

# **BIOCHEMISTRY**

## **UNIT 4 NOTES**

### **IMPORTANT TOPICS**




## BIOSYNTHESIS OF PURINE NUCLEOTIDES

There are mainly two methods of biosynthesis of Purines :

- ① De Novo Pathway
- ② Salvage Pathway

### De Novo Pathway

- In denovo pathway purine nucleotides are synthesized not free purine bases
- Ribose- 5- Phosphate used as precursor in de novo pathway :



Continues .....

5- Aminoimidazole ribosyl - 5- Phosphate

$\text{CO}_2 \rightarrow$  Carboxylase

5 - Aminoimidazole carboxylate ribosyl - 5- Phosphate

Aspartate + ATP  
ADP + Pi Synthase

5 - Aminoimidazole 4 - Succinyl  
Carboxamide ribosyl - 5- Phosphate

Fumarate  $\leftarrow$  Adenosuccinatelyase

5- Aminoimidazole 4 - carboxamide  
ribosyl - 5 Phosphate

$\text{N}^{10}$ -Methenyl THF  
THF Formyl transferase

5- Formylaminoimidazole 4- carboxamide  
ribosyl - 5- Phosphate

$\text{H}_2\text{O} \leftarrow$  Cyclohydrolase

Inosine Monophosphate

Aspartate + GTP  
GDP + Pi

Adenylsuccinate Synthase

Adenylsuccinate

Fumarate  $\leftarrow$  Adenylsuccinase

Adenosine Monophosphate  
(AMP)

$\text{NAD}^+$   
 $\text{NADH}$

IMP Dehydrogenase

Xanthosine Monophosphate




Glutamine  
Glutamate

Guanosine phosphate  
synthase

Guanosine Monophosphate  
(GMP)

## Salvage Pathway For Purines

- This pathway provides purine nucleotide for the tissues which are incapable of their biosynthesis by denovo pathway.
- example : human brain has low levels of PRPP amidotransferase & also RBCs cannot synthesize 5-phosphoribosylamine.
- Now these tissues mainly depends on salvage pathway for purine nucleotide synthesis.
- This is much simpler & required less energy.





## BIOSYNTHESIS OF PYRIMIDINE NUCLEOTIDES

Biosynthesis of pyrimidine nucleotides occurs by:


- ① De Novo Pathway
- ② Salvage Pathway

### De Novo Pathway


- The pyrimidine nucleotides are:

- ① Cytidine Monophosphate
- ② Uridine Monophosphate
- ③ Thymidine Monophosphate


- Unlike the synthesis of purine nucleotide, pyrimidine ring is made first & then attached to ribose phosphate, which is donated by PRPP.



Continues .....



## Salvage Pathway for Pyrimidines



## STRUCTURE OF DNA & RNA

### DNA

- The word DNA stands for deoxyribonucleic acid.
- It contains Deoxyribose sugar.
- DNA is made up of monomeric units called nucleotides, hence it is polymer of nucleotide or polynucleotide.


### Components

- A DNA molecule consist of double stranded nucleic acid chain which is linked via hydrogen bonds with base pairs projecting inwards.
- The hydrogen bonds attach only to specific base pairs depending on the structure of four bases.
- The Nitrogenous bases in DNA are :
  - ① Adenine
  - ② Guanine
  - ③ Thymine
  - ④ Cytosine
- The sugar phosphate group attaches in an alternate arrangement.
- The deoxyribose sugars binds with phosphodiester bonds.
- Some common types of DNA are :
  - ① Z - DNA
  - ② B - DNA
  - ③ A - DNA
  - ④ C - DNA
  - ⑤ D - DNA



## DNA Double Helix

- The double helical structure of DNA was proposed by James Watson and Francis Crick in 1953 & also got nobel prize in 1962 .
- The two strands of DNA are antiparallel to each other.



## Functions Of DNA

- DNA serves as genetic material in living being.
- DNA carry specific information to an individual.
- DNA molecules provide genetic information which implies to characteristic feature of a living organism.
- Information of all cellular protein synthesis is carried by DNA.
- The parent transfer its DNA to the offspring , so the information moves from one generation to another.
- DNA molecule has the capability of replication & transcription.

## RNA

- The word RNA stands for Ribonucleic Acid.
- It contains Ribose sugar.
- Structure of RNA is similar to DNA it is also a polymer of nucleotides.
- RNA in a cell is present in amount 10 times more than that of DNA, because RNA performs large no. of cellular functions.

### Components

A RNA molecule consist of single stranded nucleic acid chain

The nitrogenous bases in RNA are :

- Adenine
- Guanine
- Cytosine
- Uracil

### TYPES OF RNA

The three major types of RNA are :

- ① Messenger RNA
- ② Transfer RNA
- ③ Ribosomal RNA

### Messenger RNA



- Its cell composition is about 5-10%.
- It is synthesized in nucleus.
- mRNA has high molecular weight with a short life.
- mRNA of eukaryotes are more stable than prokaryotes.

## Transfer RNA

- Transfer RNA contains about 71-80 nucleotides.
- Its cell composition is about 10-20%.
- The structure of tRNA, resembles that of a clover leaf.
- tRNA contains mainly four arms with a base pair

## Ribosomal RNA

- Its cell composition is about 50-80%.
- They are the factory of protein synthesis.
- The rRNA has two subunits 50s & 30s, these subunits together form 70s ribosome



## Functions

- RNA molecules are less stable than DNA molecules. In some organism RNA molecules also carry genetic information.
- The t-RNA transport specific amino acids to ribosomes.
- The r-RNA provides structural framework to ribosomes.
- The m-RNA carries genetic information from nucleus to ribosomes.



## DNA REPLICATION

- It is the process in which DNA makes multiple copies of itself.
- It is a biological polymerisation, that proceeds in the sequence of initiation, elongation & termination.
- It is an enzyme catalysed reaction.
- DNA Polymerase is the main enzyme in the replication process.

### Models Of DNA Replication

There are basically 3 models of DNA replication :

- ① Conservative Model
- ② Dispersive Model
- ③ Semiconservative Model

### Steps of DNA Replication

There are mainly 3 steps involved in the process of DNA replication :

- ① Initiation
- ② Elongation
- ③ Termination

### INITIATION

- DNA replication demands a high degree of accuracy because even a minute mistake would result in mutations. Thus, replication cannot initiate randomly at any point in DNA.
- To begin the process of replication there is a particular region called origin of replication, this is the point where the replication originates.

Continues .....




- Replication begins with the spotting of this origin followed by the unwinding of two DNA strands.
- Unzipping of DNA strands in their entire length is not feasible due to high energy input. Hence, first, a replication fork is created catalysed by helicase enzyme that unzips the DNA strands.

### ELONGATION

- As the strands are separated, the polymerase enzymes start synthesizing the complementary sequence in each of the strands.
- The parental strands will act as a template for newly synthesising daughter strands.
- It is to be noted that elongation is unidirectional i.e. DNA is always polymerised only in the 5' to 3' direction.
- In one strand it is continuous, called continuous replication while in other strand it is discontinuous, called discontinuous replication.
- They occur as fragments called Okazaki fragments.
- The enzyme called DNA ligase joined them later.

### TERMINATION

- Termination of replication occurs in different ways in different organisms.
- An enzyme, called exonuclease removes all the RNA primers from original strand after the formation of continuous & discontinuous strands.



### ENZYMES INVOLVED IN DNA REPLICATION

DNA Replication is a highly enzyme dependent process. There are so many enzymes involved in DNA Replication as follows :

#### ① DNA Polymerase

- It helps in polymerisation, catalyses & regularises the whole process of DNA replication with support of other enzymes.
- DNA polymerase is of three types :
  - (i) DNA Polymerase I
  - (ii) DNA Polymerase II
  - (iii) DNA Polymerase III

#### ② Helicase

Helicase is the enzyme that unzips the DNA strands by breaking the hydrogen bonds between them.

### ③ Ligase

Ligase is the enzyme which joins the okazaki fragments together of discontinuous DNA strands.

### ④ Primase

The enzyme helps in the synthesis of RNA primer.

### ⑤ Endonucleases

These produce a single stranded or double stranded cut in a DNA molecule.