

# **ANATOMY & PHYSIOLOGY – II**

## **IMPORTANT QUESTIONS**

### **UNIT 5**



# **QUESTIONS - 1**

**1 DESCRIBE IN DETAIL ABOUT ANATOMY & PHYSIOLOGY  
OF MALE & FEMALE REPRODUCTIVE SYSTEM**


# FEMALE REPRODUCTIVE SYSTEM

- The Female Reproductive System refers to a complex network of reproductive organs & structures within female body that functions together to facilitate reproduction
- It's primary role is to produce eggs (ova), provide a suitable environment for fertilization, support embryo development during pregnancy & enable childbirth.

## Classification Of Female Reproductive System

It can be classified into two parts :


- ① External Genital (Vulva)
- ② Internal Genital



## EXTERNAL GENITAL ORGANS

- The external genital organs are collectively known as Vulva
- It consists of following structure :

- ① Labia Majora
- ② Labia Minora
- ③ Clitoris
- ④ Perineum
- ⑤ Vestibular Glands



### ① Labia Majora

- These are two large folds forming the boundary of Vulva.
- They are composed of skin, fibrous tissue & fat.
- They contain a large number of sebaceous & sweat glands.
- They are also known as 'Outer Lips' or 'Greater Lips'.
- They contain two Bartholin's Glands that help in lubrication during Intercourse.

### ② Labia Minora

- These are two smaller folds of skin b/w labia majora.
- They are also known as 'Inner Lips'.
- They are thin, delicate folds of skin, typically pink or reddish brown in colour.
- They provide a protective barrier for vaginal opening, urethra & clitoris.

### ③ Clitoris

- The clitoris is a highly sensitive & erectile organ, anatomically homologous to the penis in males
- It is a small pea shaped structure, containing sensory nerve endings.
- It plays a crucial role in sexual arousal & excitement in females.

### ④ Vestibular Glands

- They present on either side of urethral & vaginal orifice.
- They secrete mucus
- They are of mainly two types:
  - (i) Lesser Vestibular Glands
  - (ii) Greater Vestibular Glands.


### ⑤ Perineum

- It is the area that extends from fourchette to the anus.

## INTERNAL GENITAL ORGANS

It mainly consist of :

- Vagina
- Cervix
- Uterus
- Fallopian Tubes
- Ovaries



### ① Vagina

- Vagina is a short tubular structure.
- It is lined by mucus membrane (stratified squamous epithelium).
- It extends from vulva to the cervix of uterus.
- It receives penis during the intercourse.
- It also serves as birth canal during childbirth.
- It serves as outlet for menstrual fluid during menstruation.
- It doesn't contain any secretory glands but yet maintains a natural moisture level due to secretion of cervix & Bartholin's glands.

### ② Cervix

- The cervix is a cylindrical or conical shaped structure that forms the lower part of uterus.
- It serves as a gateway between the uterus and the vagina.

### ③ Uterus

- The uterus, also known as womb, is a pear shaped organ located in pelvic cavity of female Reproductive System
- It plays a central role in pregnancy & childbirth.
- It is responsible for menstruation in females.
- It can be divided into 3 parts :

① Fundus

② Body

③ Cervix

- It consist of 3 layers

① Endometrium (Innermost Layer)

② Myometrium (Middle Layer)

③ Perimetrium (Outer Layer)

### ④ Fallopian Tubes

- It is also known as Uterine Tubes.
- There are 2 fallopian tubes in FRS located on either side of uterus.
- They have finger like projections, called Fimbriae.
- The primary function of fallopian tubes is to transport egg from ovary to the uterus.
- It provide a suitable environment for fertilization & transportation of egg.
- Fertilization of ovum generally takes place in the upper portion of Fallopian Tubes.

## ⑤ Ovaries

- The ovaries are small, almond shaped organ located in the lower abdomen of female reproductive system.
- They play a crucial role in reproductive & hormonal functions.
- They are located on either side of uterus.
- Inside the ovary, there are thousands of tiny sacs, called follicles. each containing an immature egg (oocyte).
- The ovaries are responsible for producing & releasing eggs (ova) through a process called ovulation.
- Ovulation typically occurs once a month during menstrual cycle.
- They also produce female sex hormones i.e., Estrogen and Progesterone.

## SECONDARY SEXUAL ORGAN

It mainly contains Breast or Mammary Glands

## ① Mammary Glands


- It is also known as Breast Glands.
- It is responsible for lactation & production of milk.
- They are composed of lobules that produce milk & ducts that carry milk to the nipple.
- The growth & development of Breast tissues is stimulated by Estrogen.
- They also play a significant role in sexual attraction.

## Functions Of Female Reproductive System

- The ovaries produce & release mature eggs (ova) through a process called Ovulation.
- Female Reproductive System provides a suitable environment for fertilization of eggs by sperm.
- The ovaries produce female sex hormones, estrogen & progesterone.
- Implantation & Prenatal growth takes place in Uterus.
- Fertilization takes place in Fallopian Tubes
- The female reproductive system includes structures such as Clitoris, labia, vagina & uterus which plays role in sexual arousal, pleasure and orgasm.
- Mammary Glands of females secrete milk after Parturition.
- It is also responsible for Menstrual Cycles in females.

# MALE REPRODUCTIVE SYSTEM

- Male Reproductive system is a complex network of organs and structures responsible for producing, storing & delivering sperm cells to fertilize a female egg.
- The study of Male Reproductive System can be subdivided into two parts
  - ① External Genital Organs
  - ② Internal Genital Organs





## Internal Genital Organs

It mainly consist of :

- Testes
- Epididymis
- Vas Deferens
- Accessory Glands
  - Seminal Vesicles
  - Prostate Gland
  - Cowper's Gland
- Ejaculatory Duct

### ① Testes

- Testes are primary Male Reproductive Organs located in the scrotum, a pouch of skin situated outside of body.
- They are present in a pair.
- They are oval shaped organ responsible for both sperm production and hormone secretion.
- Spermatogenesis required  $2-3^{\circ}$  less temperature than body temperature, that's why testes lies outside of body.
- Each testis containing about 200-300 compartment called lobules.
- Now, each lobule contains approx 1-3 Seminiferous Tubules.
- Seminiferous Tubule contains Germ cells Sertoli cells & Leydig cells (Interstitial space) & primary site of Spermatogenesis.



## ② Epididymis

- The epididymis is a coiled, tubular structure located on the posterior side of each testis in Male Reproductive System.
- It is the site for sperm maturation (gaining of ability to swim) the process usually takes about 2-4 weeks.
- It also stores mature sperm cells until ejaculation.
- It further continues as Vas Deferens

## ③ Vas Deference

- The Vas Deference, also known as Ductus Deferens, is a crucial component of male reproductive system
- It is a muscular tube that transports sperm from epididymis to ejaculatory duct.
- It also serves as temporary storage site for sperm.

## ④ Accessory Glands / Ducts

They mainly contains :

- Seminal Vesicles
- Prostate Gland
- Cowper's Gland
- Ejaculatory Duct

## ① Seminal Vessides

- The seminal vessides are two small fibromuscular pouches situated near the vas deference.
- Seminal Vessides joints with vas deference to form Ejaculatory Duct.
- They secretes seminal fluid, that constitutes about 60-70% part of total volume of semen.
- Seminal Fluid mainly contains
  - Fructose
  - Prostaglandin
  - Citric Acid
  - Seminal Plasma Proteins

## ② Prostate Gland

- The prostate gland is a small, walnut shaped Organ located below the bladder.
- The prostate gland plays a crucial role in both reproductive and urinary functions.
- Prostate Gland surrounds urethra and its smooth muscle contractions helps to control the flow of urine from bladder through Urethra.
- It secretes thin, milky fluid i.e. Prostatic Fluid that constitutes about 20-30% part of semen.
- Prostatic Fluid mainly contains :
  - Citric Acid
  - Bicarbonate
  - Zinc
  - Enzymes

### © Cowper's Gland

- The Cowper's gland, also known as Bulbourethral Glands, are two small glands located near the base of penis, just below prostate gland in males.
- It helps to lubricate the urethra & tip of penis, which can reduce friction during intercourse.

### ④ Ejaculatory Ducts

- Ejaculatory Ducts are tube like structure formed by union of vas deferens & seminal vesicle.
- It transports sperm from vas deference & seminal fluid from seminal vesicle into urethra

## External Genital Organs

It mainly consist of :

- Penis
- Scrotum

### ① Penis

- Penis is a complex organ in males with both reproductive & urinary functions :
- It consist of following parts :



• Shaft : It is long cylindrical part of penis extends from base to tip

• Glans Penis : It is the enlarged tip of penis

• Urethra : A tube that runs through the penis & serves as passageway for urine & semen exit .

• Corpora Cavernosa : These are two parallel columns of erectile tissues that fills with blood during sexual arousal , causing penis erection .

• Corpus Spongiosum : It is single column of erectile tissue that contains urethra & lies below corpora cavernosa .



## ② Scrotum

- Scrotum is a sac like structure that hangs outside the body & holds the testes.
- It maintains the temperature of  $34^{\circ}\text{C}$  required for sperm production.
- It provides a protective environment for testicles.
- Scrotum is divided into two compartment by a septum

## FUNCTION OF MALE REPRODUCTIVE SYSTEM

- Testis produces sperm & testosterone.
- Epididymis stores & matures sperm.
- Penis places sperm inside body of female.
- Scrotum maintains a lower temperature for spermatogenesis
- Penis also helps in urine excretion.

## **QUESTIONS - 3**

**3 DESCRIBE VARIOUS PHASES OF MENSTRUAL CYCLE  
IN DETAILS**

# MENSTRUAL CYCLE

- The menstrual cycle is a monthly process that prepares the female body for pregnancy.
- Menstrual Cycle is defined as a cyclic event, occurring regularly in females every 26-30 days throughout the childbearing period.
- It typically lasts about 28 days but can vary from 21 to 35 days.
- It starts at the age of 12-15 years, which marks the onset of puberty.
- First Menstrual Cycle : Menarche
- Last Menstrual Cycle : Menopause

## Phases Of Menstrual Cycle

Menstrual Cycle mainly consist of following 4 phases:

- Menstrual Phase
- Follicular Phase
- Ovulatory Phase
- Luteal Phase

### ① Menstrual Phase

- Duration : Approximately days 1-5.
- Description : This phase marks the beginning of menstrual cycle, characterized by shedding of uterine lining (endometrium) if there is no pregnancy. This results in menstrual bleeding through vagina.
- Hormonal Changes : Estrogen & Progesterone levels are low.

## ② Follicular Phase

- **Duration :** Begins on first day of menstruation & lasts until ovulation, roughly days 1-14.
- **Description :** Follicular Phase involves maturation of ovarian follicles, each containing an egg.  
The Follicle Stimulating Hormone (FSH) stimulates growth of these ~~hormones~~ follicles. One dominant follicle will mature fully.
- **Hormonal Changes :** Estrogen level rises, stimulating the growth of endometrial lining in preparation for a potential implantation.

## ③ Ovulatory Phase

- **Duration :** Around day 14 but it can vary
- **Description :** Ovulation is the release of mature egg from the dominant follicle in the ovary.  
Both LH & FSH attain a peak level in the middle of cycle about 14<sup>th</sup> day  
The egg then travels down the fallopian tube, where it may be fertilized by sperm.
- **Hormonal Changes :** A surge in Luteinizing hormone (LH) triggers ovulation. Estrogen levels peak just before this surge.

#### ④ Luteal Phase

- Duration : Approximately Days 15-28
- Description : After ovulation, the ruptured follicle transforms into the corpus luteum, which secretes progesterone & some estrogen. This phase prepares the uterine lining for a possible implantation of a fertilized egg.
- Hormonal Changes : Progesterone level increases to maintain the endothelial lining. If pregnancy does not occur, corpus luteum breaks down, leading to a decrease in Progesterone levels.

## **QUESTIONS - 2**

**2 DESCRIBE THE PROCESS OF SPERMATOGENESIS**

# SPERMATOGENESIS

- Spermatogenesis is the process through which sperm cells are produced in the testes.
- Spermatogenesis takes place at temperature about  $3^{\circ}\text{C}$  below normal body temperature.
- Spermatozoa (Sperm) are produced in germ cells of seminiferous tubules of testes & matures in epididymis.

## Stages Of Spermatogenesis


Spermatogenesis takes place in two steps :

- ① Formation Of Spermatid
- ② Spermiogenesis

## Formation Of Spermatid


It further involves 3 phases :

- ① Multiplication Phase
- ② Growth Phase
- ③ Maturation Phase



## ② Spermiogenesis

- Transformation or differentiation of spermatzoa + spermatid into spermatozoa is known as Spermiogenesis.
- It is the final stage of spermatogenesis.
- Here's a detailed breakdown of spermiogenesis :
  - Condensation of Nucleus
  - Pointed, Oval, Flat
  - Formation of Acrosome
  - Loss of Excess Cytoplasm
  - Mitochondria becomes spiral
  - Maturation & Release



## **QUESTIONS - 6**

**6 DESCRIBE THE PROCESS OF OOGENESIS**

# Oogenesis

- Oogenesis is defined as process of formation of female gametes.
- Oogenesis begins in the embryonic stage.
- The process of oogenesis occurs in the follicular cells (Follicles)
- 6-7 weeks - starts oogenesis
- 20 Weeks - 6-7 million follicles
- Birth → About 2 million follicles.
- Puberty - 60,000-80,000 Primary follicles.

## Stages Of Oogenesis

It involves following key stages

- Oogonium formation
- Primary Oocyte formation
- Completion of Meiosis - I
- Secondary Oocyte & Meiosis II
- Fertilization & Meiosis II completion

### ① Oogonium Formation

During fetal development, primordial germ cells migrate to the ovaries & differentiate into oogonia, which are diploid cells.

## ② Primary Oocyte Formation

- Oogonia enter meiosis & become primary oocyte. They start meiosis - I but suspended at prophase - I until puberty.

## ③ Completion Of Meiosis - I

- At puberty, during each menstrual cycle, a primary oocyte resumes meiosis & completes first meiotic division.
- This division produces a secondary oocyte & a first polar body, which typically degenerates.

## ④ Secondary Oocyte & Meiosis - II

- The secondary oocyte begins the second meiotic division but arrests at metaphase - II, it will only complete this division if fertilization occurs.

## ⑤ Fertilization & Meiosis - II Completion

If fertilization occurs, secondary oocyte completes Meiosis - II yielding a mature ovum & secondary polar body.

Primordial Germ Cells



Oogonium




Primary Oocyte

(Suspended at Prophase of Meiosis I)

Embryonic Stage


(2n)



(First Polar body)



(n)



Secondary Oocyte

(Arrested at Metaphase II of Meiosis II)

Ovulation

Case - I



Case - II



(Second Polar body)



Fertilized Egg (Ovum)

## **QUESTIONS - 4**

**4 DESCRIBE VARIOUS PHASES OF PREGNANCY WITH DETAILED EXPLANATION ON FERTILIZATION**

# PREGNANCY

- Pregnancy occurs when sperm fertilizes the egg, and zygote is formed.
- The zygote get implanted in the wall of uterus which needs nourishment & care.
- The zygote develops into embryo which further develops into baby.
- After implantation takes place, organ development starts.
- All these steps leads to development of baby & takes about nine month.

## Stages Of Pregnancy

- Fertilization Of Ovum
- Implantation
- Development of Placenta & Embryo
- Gestation Period
- Parturition

### ① Fertilization

- Fertilization refers to fusion of male & female gametes (sperm & ovum) to form zygote.
- After sexual intercourse, semen is ejaculated in vagina, the sperms travel through the vagina & uterus to reach Fallopian Tube.

- Movement of sperm through uterus is facilitated by Antiperistalsis contraction of Uterine muscles.
- Uterine contractions are induced by oxytocin, which is secreted by posterior pituitary.
- Among 200-300 millions of sperm enters into female genital tract, only a few thousands reach the spot near the ovum. Among these thousands only 1 succeeds in fertilizing the ovum.
- The penetration of ovum by sperm is facilitated by Hyaluronidase and Proteolytic Enzymes present in acrosome of sperm.
- Immediately after fertilization, ovum (which is in secondary oocyte stage) divides into matured ovum & second polar body.
- The sperm & egg nuclei merge together to form a single diploid nucleus.
- The combination of genetic material from both parents results in the formation of zygote, which then begins to divide & develop into an Embryo.

## ② Implantation

- Implantation is the process by which the fertilized ovum called Zygote implants (gets attached) in the endometrial lining of uterus.
- Zygote takes 3-5 days to reach the uterine cavity from Fallopian Tube.

### ③ Development of Placenta & Embryo

- After implantation, a placenta is formed in uterine muscles to support fetal development throughout pregnancy.
- When implantation occurs, there is further increase in the thickness of endometrium because of continuous secretion of Progesterone from Corpus Luteum.
- An Embryo is the early stage of development of a multicellular organism.

### ④ Gestation Period

- It refers to pregnancy period & measured in weeks.
- A normal pregnancy can range from 38-42 weeks.
- Infants born before 37 weeks are considered premature.

### ⑤ Parturition

- Parturition is the delivery of fetus from mother's body.
- It occurs at end of pregnancy.
- The process by which delivery of fetus occurs is called labor.

# **QUESTIONS - 5**


## **5 DESCRIBE GENES / CHROMOSOMES**

# CHROMOSOMES

- Chromosomes are thread like structures composed of DNA & Proteins that are found in the nucleus of eukaryotic cells.
- They carry genetic information necessary for growth, development & functioning of all living organism.

## Structure

- Chromosomes primarily composed of Deoxyribonucleic Acid (DNA) which contains genetic blueprint of an organism.
- Along with DNA, DNA contains Histone proteins that helps in organizing and compacting DNA into more manageable structure.



Replicated  
Chromosome

## Number

Except reproductive cells (sperm & ovum) all the human cells have 46 chromosomes arranged in 23 pairs, including 22 pair of Autosomes and 1 pair of sex chromosomes.

## Types Of Chromosomes

There are mainly two types of chromosomes :

- ① Autosomes
- ② Sex Chromosomes

### ① Autosomes

- These are chromosomes that are not involved in determining the sex of an organism ,
- Humans have 22 pair of autosomes

### ② Sex Chromosomes

- These chromosomes determines the sex of an individual
- Females - XX
- Males - XY

## Functions Of Chromosomes

- Chromosomes carry genes , the units of heredity that encode specific proteins .
- During cell division , chromosomes ensures that DNA is accurately copied & distributed to daughter cells .

# THANK YOU

FOR CHOOSING IMPERFECT PHARMACY AS YOUR STUDY PARTNER



## JOIN US ON:



**IMPERFECT PHARMACY**



**IMPERFECT PHARMACY**



**@IMPERFECTPHARMA**



**IMPERFECT PHARMACY**



**@IMPERFECTPHARMACY**



**IMPERFECT PHARMACY**



**IMPERFECTPHARMACY@GMAIL.COM**